

Effect of Copper Dust on Photosynthesis Pigments Concentration in Plants Species

Mohnish Pichhode, Kumar Nikhil

Abstract— Effect of copper concentration was studied in the plants within the Malanjkhanda Copper Mining Project (MCP), Balaghat, Madhya Pradesh (India). Copper affects vegetation were determined through biochemical parameter (paper chromatography) in the tree species like *Tectona grandis*, *Mangifera indica*, *Butia monosperma* and *Madhuca indica* which were exposed to different copper dust concentration load for short duration. Variations in biochemical parameter like different pigments in the leaves were separated by paper chromatography found significant in difference. This variation can be used as indicator of copper concentration for early diagnosis of stress as a maker for physiological damage to plants prior to the onset of visible injury symptoms. The present study reflects the elevated concentration of copper in the associated plant species within the Malanjkhanda Copper Mining Site.

I. INTRODUCTION

Mining has been one of the most common activities since ancient times and continues to remain so in the modern world. Mining is an important part of our economy. Minerals extracted raw from earth, are processed to yield basic substances such as metals, chemicals, building materials, fuels, fertilizers etc. industrial society could not exist without these essential commodities. A combined total of about 1150 million tones of heavy metals (copper, lead, cobalt, zinc, cadmium and chromium) have been mined by man since the Stone Age. It is further estimated that an annual output of 14 million tones of heavy metals is being mined with annual growth of 3.4% (Matagi *et al.*, 1998). The continued advancement in industrialization and the ever increasing demand for energy resources and minerals, have spurt in mining activities, bringing in its wake imbalances in ecological equilibrium and many environmental hazards (Wu *et al.*, 2007; Vamerali *et al.*, 2010).

Mining activities such as crushing, grinding, washing, smelting, and all other processes used to extract, concentrate generate waste products such as mine overburden and mine's waste soils. As a result, very significant volumes of wastes have been deposited on soil and wild plants and animals are exposed to elements contained in the residue. People living near these sites are also exposed through wind and soil erosion. The direct effect will be loss of cultivated land, forest

Manuscript received Feb 14, 2015

Mohnish Pichhode, Department of Biotechnology, Sardar Patel College of Technology, Balaghat, India,

Dr. Kumar Nikhil, Principal Scientist, Environmental management Group CSIR-Central Inst. Of Mining & Fuel Research, Dhanbad, India

or grazing land, and the overall loss of production. The indirect effects include air and water pollution and water pollution and siltation of water body. This will eventually lead to loss of biodiversity, amenity, and economic wealth (Yang *et al.*, 2002; Wong, 2003). The management of these waste materials is an important issue for mining industries worldwide.

Copper is essential as micronutrient for microorganisms, plants and animals while others have no known biological function (Welch, 1995). All heavy metals at high concentrations have strong toxic effects and regarded as environmental pollutants (Nedelkoska and Doran, 2000; chehregani *et al.*, 2005). Contaminated soil negatively affects crop growth because of interference of phytotoxic contamination with metabolic processes and leading to plant death (Pal and Rai, 2010).

The present research aimed to investigate the influence of mining activity on the concentration of heavy metal as copper in the soil and vegetation surrounding copper mining sites at Malanjkhanda located in Balaghat, Madhya Pradesh, India.

II. MATERIALS AND METHOD

A. Sampling Site

The study area Malanjkhanda is located in the Balaghat district of Madhya Pradesh 250 km northeast of Nagpur (MH). The Malanjkhanda town is connected by major arterial roads, some of which are NH 16 and other district road together constitute the primary network of major roads. Malanjkhanda referred to as MCP (Malanjkhanda Copper Project) is an open-pit copper mine in India, located near the town of Malanjkhanda, 90 kilometers (56 miles) northeast of Balaghat in Madhya Pradesh at an altitude of 576 MRL. It falls in the tehsil of Baihar, which is 22 km from the project site; on the way to Balaghat headquarter. Malanjkhanda open-pit copper mine is largest metal mine in India. The special thematic mapping during 1991 & 1994 by Jaggi *et al.*, (GSI) has brought out the regional geological set-up. Malanjkhanda copper belt lies between latitude 22°00'59"-22°02'24" and longitude 80°41'51"- 80°42'38" and from a part of Survey of India toposheet no. 64 B/12 (Fig.1).

Malanjkhanda copper project was established in 1982. Initial project has been set up by Hindustan Copper Limited (HCL) to exploit the copper ore through an open-pit mine. Geological survey of India took systematic geological exploration as this deposit during 1969. Mining lease of the ore was granted to HCL during 1973. With advancement of time this project was enhanced with viable operational developments. In fact it is largest open cast mining in Asia. This area is the result of many cycles of erosion on the gneiss base during the Amgaon orogeny.




Figure 1: Geographical location of study site Malanjkhanda, District- Balaghat, (M.P.), India.

B. Plant Sampling and Analysis

Firstly the leaves were plucked at chest level height of the ten plants of four species sampled nearby road side from where the ore were carried from mining to storage site with the trafficking of dumpers. The same species were sampled also from deep forest far away from the mining site having no affect of copper or any other dust. The plant species taken for this study were *Tectona grandis*, *Mangifera indica*, *Butia monosperma* and *Madhuca indica*. The leaves were plucked and collected in a full newspaper assuring not to loose any part of dust from leaf. Immediately leaf were washed with distilled water, filtered and dust were collected after drying filter paper. Further, the washed leaves were crushed in mortar and pestle and mixed with different solvents and for the separation of different pigments loaded on whatman filter paper no. 1 for paper chromatography assembly (Fig.2).

The Rf (retardation factor) value of the pigments were found with a particular solvents for the four planted species were shown in the **Table.1 and 2** without dust and with dust affected plant species leaves respectively.

III. RESULTS AND DISCUSSION

A. Deposition of Dust

Plant species leaves were collected and studied for the dust deposition at mining sites. Dust was collected through washing the leaves and evaporating the water. The dust an average was found $\pm 1.344, \pm 0.783, \pm 0.839, \pm 0.731$ gm from *Tectona grandis* (Teak), *Mangifera indica* (Mango), *Butia monosperma* (Palash), *Madhuca indica* (Mahua) respectively of an average of ten plants of each four species.

B. Separation of Pigments

The leaves affected by dust were studied for the carotenoid, xanthophylls, chlorophyll a and chlorophyll b content in all the four planted species at mining and as well as forest as control sites. Significant results were found while comparing the results of different pigmentation in the four plant species leaves in control and mining affected site.

The Rf value (Retardation factor) of carotinioid, xanthophylls, chlorophyll a, chlorophyll b pigments of control as well as copper mining affected plants were assessed.






Figure 2: Shows the separation of pigments on whatman filter paper

The data were statistically analyzed for standard deviation of the Rf (retardation factor) value within all the four plants species in different photosynthesis pigments in leaves with dust and without dust .

Plant pigments	Mean (X)	Mode (Z)	Median (M)	Standard deviation (σ)
<i>Tectona grandis</i>				
Carotenoid	0.955	0.969	0.955	0.014
Xanthophylls	0.932	0.939	0.932	0.0065
Chlorophyll a	0.872	0.878	0.872	0.0055
Chlorophyll b	0.806	0.818	0.806	0.0130
<i>Mangifera indica</i>				
Carotenoid	0.977	0.984	0.977	0.0065
Xanthophylls	0.919	0.954	0.919	0.035
Chlorophyll a	0.882	0.909	0.882	0.027
Chlorophyll b	0.785	0.818	0.785	0.0325
<i>Butea monosperma</i>				
Carotenoid	0.956	0.970	0.956	0.014
Xanthophylls	0.927	0.955	0.927	0.0275
Chlorophyll a	0.876	0.910	0.876	0.034
Chlorophyll b	0.781	0.805	0.781	0.0229

<i>Madhuca indica</i>				
Carotenoid	0.955	0.955	0.955	000
Xanthophylls	0.903	0.911	0.903	0.008
Chlorophyll a	0.843	0.852	0.843	0.0085
Chlorophyll b	0.792	0.791	0.792	0.0015
Mix				
Carotenoid	0.964	0.971	0.964	0.007
Xanthophylls	0.9005	0.901	0.9005	0.0005
Chlorophyll a	0.808	0.816	0.808	0.008
Chlorophyll b	0.758	0.760	0.758	0.001

Table 1: Rf value of different pigments for the plant species leaves without copper dust




Graph 1: Rf value of different pigments in plant species leaves without copper dust

Plant pigments	Mean (X)	Mode (Z)	Median (M)	Standard deviation (σ)
<i>Tectona grandis</i>				
Carotenoid	000	000	000	000
Xanthophylls	0.895	0.9	0.895	0.005
Chlorophyll a	0.846	0.857	0.846	0.011
Chlorophyll b	0.727	0.728	0.727	0.001
<i>Mangifera indica</i>				
Carotenoid	0.767	0.771	0.767	0.004
Xanthophylls	0.711	0.722	0.767	0.011
Chlorophyll a	0.661	0.666	0.661	0.0045
Chlorophyll b	0.626	0.642	0.626	0.0219
<i>Butea monosperma</i>				
Carotenoid	0.846	0.847	0.846	0.001
Xanthophylls	0.748	0.760	0.748	0.012
Chlorophyll a	0.72	0.732	0.72	0.0114
Chlorophyll b	0.664	0.676	0.664	0.012
<i>Madhuca indica</i>				
Carotenoid	0.875	0.887	0.875	0.0115
Xanthophylls	0.841	0.845	0.841	0.004
Chlorophyll a	0.778	0.783	0.778	0.0045

Chlorophyll b	000	000	000	000
Mix				
Carotenoid	0.863	0.864	0.863	0.0007
Xanthophylls	0.836	0.837	0.836	0.001
Chlorophyll a	0.802	0.810	0.802	0.008
Chlorophyll b	0.679	0.684	0.679	0.0045

Table 2: Rf value of different pigment for the plant species leaves affected by copper dust



Graph 2: Rf value of different pigment in plant species leaves affected by copper dust

Graph.1 & 2 clearly reveals that deposition of copper dust has substantially lowered down the four pigments in all the four plant species studied having almost same age from both the sites.

Air pollutants, fly ash and dust emissions have a profound impact on the concentration of different photosynthetic pigments. Polluted and dusted leaf surface is responsible for reduced photosynthetic and thereby causing reduction in chlorophyll content (Kalyani, and Singaracharya, 1995). The similar impact of air pollutants in the concentration of chlorophyll contents have been reported by a number of other works (Nikova and Dushkova; Rabe and Kareeb, 1979; Dubey and Pawar, 1985; Shah, et., al.,1989; Saxena, 1991; Swami, et.al., 2004 and Tripathi and Gautam, 2007). The photosynthetic pigments are the most likely to be damaged by air pollution.

Chlorophyll pigments exist in highly organized state, and under stress they may undergo several photochemical reactions such as oxidation, reduction, pheophytinisation and reversible bleaching (Puckett, et.al.). Hence any alteration in chlorophyll concentration may change the morphological, physiological and biochemical behaviour of the plant. Air pollution-induced degradation in photosynthetic pigments was also observed by a number of workers (Bansal, 1988; Singh, et al., 1990 and Sandelius, et.al., 1995).

In all the four plants species chlorophyll a and chlorophyll b content were reduced significantly at polluted site.

CONCLUSION

In the present study, effect of copper dust was seen on the concentration of four photosynthesis pigments in the leaves of four plants in and around mining and forest areas. The plant species were *Tectona grandis*, *Mangifera indica*, *Butia*

monosperma, *Madhuca indica* plant species were studied to see the effects of road side dust such as copper dust, vehicular dust pollutants have been investigated and their effects are presented. From the studies it is concluded that the dust pollutants causes degradation in plant growth. Never the less the experiment conducted in the hypothesis of proving the significant effect of copper dust on the concentration of the pigmentation in four species leaves, which are highly sensitive and overall growth compared from same plant species at control condition species i.e., forest site.

ACKNOWLEDGEMENT

The authors are grateful to Director, CSIR-Central Institute of Mining and Fuel Research (CIMFR), Dhanbad, Jharkhand, India and Principal, Sardar Patel College of Technology, Balaghat, M.P., India for giving necessary permission for conducting in house experiment and publishing this paper as a part from the report submitted for the summer training program.

REFERENCES

- [1] Alaoui-Sosse, B., P. Genet, F. Vinit-Dunand, M.L. Taussaint, D. Epran and P.M. Badot, Effect of copper on growth in cucumber plants and its relationships with carbohydrate accumulation and changes in ion content, *Plant Science*, 2004, pp. 1213-1218.
- [2] Chatter, H., Effect of dust on the enzymatic activity in the levels of *Triticum aestivum*, *Acta Ecologica*, 1991, pp. 113-119.
- [3] Chehregani A., Malayeri B. and Goltmohammadi R., Effect of heavy metals on the developmental stages of ovules and embryonic sac in *Euphorbia cheirandenia*, *Pakistan Journal of Biological Sciences*, 2004, pp. 622-625.
- [4] Farmer Andrew M., The effects of dust on vegetation- a review, *Environmental pollution*, Elsevier Science Publishers Ltd, England, 1993, pp. 63-75.
- [5] Fikriye Kribag Zengin and sevda Kirbag, Effects of copper on chlorophyll, proline, protein, and abscisic acid level of sunflower seedlings, *Journal of Environmental Biology*, 2007, pp. 561-566.
- [6] Gareth, J., R.Alan, F. Jean and H.Graham, *Environmental science*, Harper perennial, Division of harper Collins publishers, 1192, pp. 564.
- [7] Gupta A.K., Mishra R.M., Effect of lime kinos air pollution on some plant species, *Pollution Research* 13 (1), 1994.
- [8] Kupper H.F. and Spiller M., Environmental relevance of heavy metal substituted chlorophylls using the example of water plants, *Journal of Experimental Botany*, 1996, pp. 275-280.
- [9] Lone M.I.He., Z. Stoffella P.J. and Yang X., Phytoremediation of heavy metal polluted soil and water: progress and perspectives, *Journals of Zhejiang university Science*, 2008, pp. 210-220.
- [10] Matagi S.V., Swai D. and Mugabe R., A review of heavy metal removal mechanisms in wetlands, *African Journal Tropical Hydrobiology Fish*, 1998, pp. 23-35.
- [11] Nedelkoska T.V. and Doran P.M., Hyper accumulation of Cadmium by hairy roots of *Thlaspi caerulescens*, *Biotechnology and Bioengineering*, 2000, pp. 607-615.
- [12] Nriagu J.O. and pacyna J.M., Quantitative assessment of worldwide contamination of air, water and soils by trace elements, *Nature* 333(6169), 1988, pp. 134-139.
- [13] Pal R. and Rai J.P.N., Phytochelatin: peptides involved in Heavy metal detoxification, *Applied Biochemistry and Biotechnology*, 2010, pp. 945-963.
- [14] Sheoran V., Poonia P., Trivedi S.K., Metal pollution in soil and plants near copper mining site, *International Journal of Geology, Earth and environmental Sciences*, 2011, pp. 27-34.
- [15] Tripathi A.K., Gautam Mukesh, biochemical parameters of plants as indicators of air pollution, *journal of Environmental Biology*, 2007, pp. 127-132.
- [16] Vamerali T. and Marianna B. Mosca G., Field crops for phytoremediation of metal-contaminated of metal-contaminated land: A review, *Environmental Chemistry Letters*, 2010, pp. 1-17.
- [17] Welch R.M., Micronutrient nutrition of plants, *Critical Reviews in Plant Sciences*. 1995, pp. 49-82.
- [18] Wu Q.T., Wei Z.B., Ouyang Y., Phytoextraction of metal contaminated soil by *Sedum alfredi*: effect of chelators and co-planting, *WaterAir and Soil Pollution*, 2007, pp. 131-139.
- [19] Yang X Long XX Ni WZ, Physiological and Molecular mechanisms of heavy metal uptake by hyperaccumulating plants, *Plant Nutrition and fertilization Science*, 2002, pp. 8-15.
- [20] Kalyani, V., and Singaracharya M.A., "Biomonitoring of air pollution in Warangal city (A.P.)". *Act Botanica indica*; **23**:21-23, (1995).
- [21] Nikova, D. and Dushkova, P. 919780. *Gorshostoparnska Nauka*. **3**: 3-8.
- [22] Rabe, R. and Kareeb, K. H., Enzyme activities and chlorophyll and protein content in plants as indicator of air pollution. *Environ. Pollut.* **16**:119-137, (1979).
- [23] Dubey, P. S. and Pawar, K., Air pollution and plant response- Review. In: *Perspective in environment botany* (Eds.:D. N. Rao, K. J. Ahamed. M. Yunus and S. N. Singh). Print House, Lucknow. **1**: 101-118, (1985).
- [24] Shah, F. H., Ilahi, I. and Rashid, A., Effect of cement dust on the chlorophyll contents, stomatal clogging and biomass of some selected plants. *Pakistan J. Sci. Indust. Research*. **32**(8): 542-545, (1989).
- [25] Saxena, R. M., Effect of motorway pollution on seed health of some vegetable crops. *Ind. J. Environ. Hlth.* **33**: 385-387, (1991).
- [26] Swami, A. D., Bhatt and Joshi, P. C., Effect of automobile pollution on Sal (*Shorea robusta*) and Rohini (*Mallotus philippinensis*) at Asarori, Dehradun. *Him. J. Environ. Zool.* **18**: 57-61, (2004).
- [27] Tripathi, A. K. and Gautam, M. Biochemical parameters of plants as indicators of air pollution. *J. Environ. Biol.* **28**: 127-132, (2007).
- [28] Puckett, K.J., Nieboer, E., Flora W.P., and Richardson, D.H.S. "Sulphur dioxide: Its effect on photosynthetic ¹⁴C fixation in lichens and suggested mechanism of phytotoxicity" *The New Phytologist*, **72**, pp.141-154.
- [29] Bansal, S., "Studies on the effect of certain atmospheric pollutants on fruit diseases of *Lycopersicon esculentum* Mill. Caused by *Alternaria alternate*", PhD thesis, Bhopal, Bhopal University, (1988).
- [30] Singh, N., Singh, S.N., Srivastava, K., Yunus, M., Ahmad, K.J., and Sharma S. C., et al., "Relative sensitivity and tolerance of some *Gladiolus* cultivars to sulphur dioxide", *Annals of Botany*, **65**, pp. 41-44, (1990).
- [31] Sandelius, A.S., Naslund, K., Carlson, A.S., Pleijel, H., and Sellden, G., "Exposure of spring wheat (*Triticum aestivum*) to ozone in open top chambers. Effects on acyl lipid composition and chlorophyll content of flag leaves", *The New Phytologist*, **131**, pp. 231-239, (1995).



Mohnish Pichhode, M.Sc in Biotechnology, Department of Biotechnology, Sardar Patel College of Technology, Balaghat affiliated to Rani Durgavati Vishwavidyalaya, Jabalpur, M.P done his project work at EMG, CSIR-CIMFR, Dhanbad, Jharkhand, India in the year 2013 and publishing his experimental finding which is very significant and important work helps in understanding the physiology and morphological behavior of plant species in copper mining area.



Dr. Kumar Nikhil, Ph.D in Env.Sc. & Engg., Principal Scientist at Environmental Management Group, CSIR- Central Institute of Mining and Fuel Research (CIMFR), Barwa Road, Dhanbad - 826 015, Jharkhand, India has gained more than 30 years of research experience involved in more than 60 projects in different capacity. More than 135 scientific publications on his name. Guided more than 60 students of B.Sc, M.Sc, B.Tech & M.Tech, Ph.D students in their project and research works.